

121512

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS, SHALL COME;

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

*September 15, 2004*

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/553,963  
FILING DATE: *March 18, 2004*

Certified by



Jon W Dudas

Acting Under Secretary of Commerce  
for Intellectual Property  
and Acting Director of the U.S.  
Patent and Trademark Office



13281 U.S. PTO

## PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION under 37 CFR 1.53(c).

19587 U.S. PTO  
60/53963

031804

Docket Number		50193-178	Type a plus sign (+) inside this box →	+
INVENTOR(s)/APPLICANT(s)				
LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (City and Either State or Foreign Country)	
HAMMOND FUKUDA	Milton Yasumichi	L.	Branchburg, New Jersey Oyama Toahigi, JAPAN	
TITLE OF THE INVENTION (280 characters max)				
OXAZOLIDINONE ANTIBIOTICS AND DERIVATIVES THEREOF				
CORRESPONDENCE ADDRESS				
McDERMOTT, WILL & EMERY 600 13th Street, N.W. Washington, D. C. 20005-3096 (202) 756-8000				
STATE	Washington, D. C.	ZIP CODE	20005-3096	COUNTRY USA
ENCLOSED APPLICATION PARTS (check all that apply)				
<input checked="" type="checkbox"/>	Specification	Number of pages [56]	<input type="checkbox"/>	Small Entity Statement
<input type="checkbox"/>	Drawings	Number of sheets	<input type="checkbox"/>	Other (specify):
METHOD OF PAYMENT (check one)				
<input type="checkbox"/>	A check or money order is enclosed to cover the Provisional filing fees			PROVISIONAL FILING FEE AMOUNT (\$)
<input checked="" type="checkbox"/>	The Commissioner is hereby authorized to charge filing fees and credit Deposit Account Number: 500417			\$160.00

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.



No.



Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

McDERMOTT, WILL &amp; EMERY

Judith L. Toffenetti  
Registration No. 39,048600 13<sup>th</sup> Street, N.W.  
Washington, DC 20005-3096  
(202) 756-8000 JLT:gav  
Facsimile: (202) 756-8087  
Date: March 18, 2004

# TITLE OF THE INVENTION

## Oxazolidinone Antibiotics and Derivatives Thereof


### BACKGROUND OF THE INVENTION

5 Oxazolidinones represent the first new class of antibacterials to be developed since the quinolones. The oxazolidinones are synthetic antibacterial compounds that are orally or intravenously active against problematic multidrug resistant Gram positive organisms and are not cross-resistant with other antibiotics. See Riedl et al, Recent Developments with Oxazolidinone Antibiotics, *Exp. Opin. Ther. Patents* (1999) 9(5), Ford et al., Oxazolidinones: New Antibacterial Agents, *Trends in Microbiology* 196 Vol.5, No. 5, 10 May 1997 and WO 96/35691. See also WO 03/063862, WO 01/81350, WO 01/94342, WO 03/072553, EP 0352781 and US 5,565,571 and 4,053,593.

15 This invention relates to new oxazolidinones having a cyclopropyl moiety, which are effective against aerobic and anerobic pathogens such as multi-resistant staphylococci, streptococci and enterococci, *Bacteroides* spp., *Clostridia* spp. species, as well as acid-fast organisms such as *Mycobacterium tuberculosis* and other mycobacterial species.

### SUMMARY OF THE INVENTION

20 The present invention relates to compounds of formula I:



25 its enantiomer, diastereomer, or pharmaceutically acceptable salt, hydrate or prodrug thereof wherein:

R<sub>1</sub> represents

- i) hydrogen,
- ii) NR<sub>5</sub>R<sub>6</sub>,

- iii)  $CR_7R_8R_9$ ,  $C(R)_2OR_{14}$ ,  $CH_2NHR_{14}$ ,  
 iv)  $C(=O)R_{13}$ ,  $C(=NOH)H$ ,  $C(=NOR_{13})H$ ,  $C(=NOR_{13})R_{13}$ ,  $C(=NOH)R_{13}$ ,  $C(=O)N(R_{13})_2$ ,  
 $C(=NOH)N(R_{13})_2$ ,  $NHC(=X_1)N(R_{13})_2$ ,  $(C=NH)R_7$ ,  $N(R_{13})C(=X_1)N(R_{13})_2$ ,  $COOR_{13}$ ,  
 $SO_2R_{14}$ ,  $N(R_{13})SO_2R_{14}$ ,  $N(R_{13})COR_{14}$ ,  
 5 v)  $(C_{1-6}alkyl)CN$ ,  $CN$ ,  $CH=C(R)_2$ ,  $(CH_2)_pOH$ ,  $C(=O)CHR_{13}$ ,  $C(=NR_{13})R_{13}$ ,  
 $NR_{10}C(=X_1)R_{13}$ ; or

vi)  $C_{5-10}$  heterocycle optionally substituted with 1-3 groups of  $R_7$ , which may be attached through either a carbon or a heteroatom;

10

A represents  $NR$ , O, or  $S(O)_p$ ;



- represents aryl or heteroaryl, heterocycle, heterocyclyl or heterocyclic, provided that in the case of a heteroaryl, heterocycle, heterocyclyl or heterocyclic, the cyclopropyl is  
 15 not attached to a nitrogen atom on the ring;

$R_x$  represents hydrogen or  $C_{1-6}$  alkyl;

- $R_3$  represent  
 20 i)  $NR_{13}(C=X_2)R_{12}$ ,  
 ii)  $NR_{13}(C=X_1)R_{12}$ ,  
 iii)  $NR_{13}SO_2R_{14}$ ,  
 iv)  $N(R_{13})heteroaryl$ ,  
 v)  $NR_{13}(CHR_{13})_{0-4}aryl$ ,  
 25 vi)  $NR_{13}(CHR_{13})_{0-4}heteroaryl$ ,  
 vii)  $S(CHR_{13})_{0-4}aryl$ ,  
 viii)  $S(CHR_{13})_{0-4}heteroaryl$ ,  
 ix)  $O(CHR_{13})_{0-4}aryl$ ,  
 x)  $O(CHR_{13})_{0-4}heteroaryl$ ,  
 30 xi)  $NOH(C=X_1)R_{12}$ ,  
 xii)  $-OC=N(OCOaryl) C_{1-6} alkyl$   
 xiii)  $-OC=N(OH) C_{1-6} alkyl$

xiv) C<sub>5-10</sub> heteroaryl which may be attached through either a carbon or a heteroatom; said aryl and heteroaryl optionally substituted with 1-3 groups of R<sub>7</sub>,

R<sub>4</sub>, and R<sub>4a</sub>, independently represent

- 5 i) hydrogen,
- ii) halogen,
- iii) C<sub>1-6</sub> alkoxy, or
- iv) C<sub>1-6</sub> alkyl

- 10 r and s independently are 1-3, with the provision that when (R<sub>4a</sub>)<sub>s</sub> and (R<sub>4</sub>)<sub>r</sub> are attached to an Ar or HAr ring the sum of r and s is less than or equal to 4;

R<sub>5</sub> and R<sub>6</sub> independently represent

- i) hydrogen,
- 15 ii) C<sub>1-6</sub> alkyl optionally substituted with 1-3 groups of halogen, CN, OH, C<sub>1-6</sub> alkoxy, amino, imino, hydroxyamino, alkoxyamino, C<sub>1-6</sub> acyloxy, C<sub>1-6</sub> alkylsulfenyl, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylsulfonyl, aminosulfonyl, C<sub>1-6</sub> alkylaminosulfonyl, C<sub>1-6</sub> dialkylaminosulfonyl, 4-morpholinylsulfonyl, phenyl, pyridine, 5-isoxazolyl, ethylenyloxy, or ethynyl, said phenyl and pyridine optionally substituted with 1-3
- 20 halogen, CN, OH, CF<sub>3</sub>, C<sub>1-6</sub> alkyl or C<sub>1-6</sub> alkoxy;
- iii) C<sub>1-6</sub> acyl optionally substituted with 1-3 groups of halogen, OH, SH, C<sub>1-6</sub> alkoxy, naphthalenoxy, phenoxy, amino, C<sub>1-6</sub> acylamino, hydroxylamino, alkoxyamino, C<sub>1-6</sub> acyloxy, aralkyloxy, phenyl, pyridine, C<sub>1-6</sub> alkylcarbonyl, C<sub>1-6</sub> alkylamino, C<sub>1-6</sub> dialkylamino, C<sub>1-6</sub> hydroxyacyloxy, C<sub>1-6</sub> alkylsulfenyl, phthalimido, maleimido,
- 25 succinimido, said phenoxy, phenyl and pyridine optionally substituted with 1-3 groups of halo, OH, CN, C<sub>1-6</sub> alkoxy, amino, C<sub>1-6</sub> acylamino, CF<sub>3</sub> or C<sub>1-6</sub> alkyl;
- iv) C<sub>1-6</sub> alkylsulfonyl optionally substituted with 1-3 groups of halogen, OH, C<sub>1-6</sub> alkoxy, amino, hydroxylamino, alkoxyamino, C<sub>1-6</sub> acyloxy, or phenyl; said phenyl optionally substituted with 1-3 groups of halo, OH, C<sub>1-6</sub> alkoxy, amino, C<sub>1-6</sub> acylamino, CF<sub>3</sub> or C<sub>1-6</sub> alkyl;
- 30 v) arylsulfonyl optionally substituted with 1-3 of halogen, C<sub>1-6</sub> alkoxy, OH or C<sub>1-6</sub> alkyl;

- vi) C1-6 alkoxy carbonyl optionally substituted with 1-3 of halogen, OH, C1-6 alkoxy, C1-6 acyloxy, or phenyl, said phenyl optionally substituted with 1-3 groups of halo, OH, C1-6 alkoxy, amino, C1-6 acylamino, CF<sub>3</sub> or C1-6 alkyl;
- vii) aminocarbonyl, C1-6 alkylaminocarbonyl or C1-6 dialkylaminocarbonyl, said alkyl groups optionally substituted with 1-3 groups of halogen, OH, C1-6 alkoxy or phenyl
- 5 viii) five to six membered heterocycles optionally substituted with 1-3 groups of halogen, OH, CN, amino, C1-6 acylamino, C1-6 alkylsulfonylamino, C1-6 alkoxy carbonylamino, C1-6 alkoxy, C1-6 acyloxy or C1-6 alkyl, said alkyl optionally substituted with 1-3 groups of halogen, or C1-6 alkoxy;
- 10 ix) C3-6 cycloalkyl carbonyl optionally substituted with 1-3 groups of halogen, OH, C1-6 alkoxy or CN;
- x) benzoyl optionally substituted with 1-3 groups of halogen, OH, C1-6 alkoxy, C1-6 alkyl, CF<sub>3</sub>, C1-6 alkanoyl, amino or C1-6 acylamino;
- xi) pyrrolyl carbonyl optionally substituted with 1-3 of C1-6 alkyl;
- 15 xii) C1-2 acyloxyacetyl where the acyl is optionally substituted with amino, C1-6 alkylamino, C1-6 dialkylamino, 4-morpholino, 4-aminophenyl, 4-(dialkylamino)phenyl, 4-(glycylamino)phenyl; or

R<sub>5</sub> and R<sub>6</sub> taken together with any intervening atoms can form a 3 to 7 membered heterocyclic ring containing carbon atoms and 1-2 heteroatoms independently chosen from  
 20 O, S, SO, SO<sub>2</sub>, N, or NR<sub>8</sub>;

R<sub>7</sub> represent

- i) hydrogen, halogen, CN, CO<sub>2</sub>R, CON(R)<sub>2</sub>, CHO, CH<sub>2</sub>NHAc, C(=NOR), OH, C1-6 alkoxy, C1-6 alkyl, alkenyl, hydroxy C1-6 alkyl, (CH<sub>2</sub>)<sub>1-3</sub>NHC(O)C1-6 alkyl, (CH<sub>2</sub>)<sub>1-3</sub>N(C1-6 alkyl)<sub>2</sub>
- 25 ii) (CH<sub>2</sub>)<sub>n</sub>amino, (CH<sub>2</sub>)<sub>n</sub>C1-6 alkylamino, C1-6 dialkylamino, hydroxylamino or C1-2 alkoxyamino all of which can be optionally substituted on the nitrogen with C1-6 acyl, C1-6 alkylsulfonyl or C1-6 alkoxy carbonyl, said acyl and alkylsulfonyl optionally substituted with 1-2 of halogen or OH;

30

R<sub>8</sub> and R<sub>9</sub> independently represents

- i) H, CN,
- ii) C1-6 alkyl optionally substituted with 1-3 halogen, CN, OH, C1-6 alkoxy, C1-6 acyloxy, or amino,

iii) phenyl optionally substituted with 1-3 groups of halogen, OH, C<sub>1-6</sub> alkoxy; or

R<sub>7</sub> and R<sub>8</sub> taken together can form a 3-7 membered carbon ring optionally interrupted with 1-2 heteroatoms chosen from O, S, SO, SO<sub>2</sub>, NH, and NR<sub>8</sub>;

5

X<sub>1</sub> represents O, S or NR<sub>13</sub>, NCN, NCO<sub>2</sub>R<sub>16</sub>, or NSO<sub>2</sub>R<sub>14</sub>

X<sub>2</sub> represents O, S, NH or NSO<sub>2</sub>R<sub>14</sub>;

10 R<sub>10</sub> represents hydrogen, C<sub>1-6</sub> alkyl or CO<sub>2</sub>R<sub>15</sub>;

R<sub>12</sub> represents hydrogen, C<sub>1-6</sub> alkyl, NH<sub>2</sub>, OR, CHF<sub>2</sub>, CHCl<sub>2</sub>, CR<sub>2</sub>Cl, (CH<sub>2</sub>)<sub>n</sub>SR, (CH<sub>2</sub>)<sub>n</sub>CN, (CH<sub>2</sub>)<sub>n</sub>SO<sub>2</sub>R, (CH<sub>2</sub>)<sub>n</sub>S(O)R, C<sub>1-6</sub> alkylamino, C<sub>5-10</sub> heteroaryl or C<sub>1-6</sub> dialkylamino, where said alkyl may be substituted with 1-3 groups of halo, CN, OH or C<sub>1-6</sub> alkoxy, said

15 heteroaryl optionally substituted with 1-3 groups of R<sub>7</sub>;

Each R<sub>13</sub> represents independently hydrogen, C<sub>1-6</sub> alkyl, C<sub>6-10</sub> aryl, NR<sub>5</sub>R<sub>6</sub>, SR<sub>8</sub>, S(O)R<sub>8</sub>, S(O)<sub>2</sub>R<sub>8</sub>, CN, OH, C<sub>1-6</sub> alkylS(O)R, C<sub>1-6</sub> alkoxy carbonyl, hydroxycarbonyl, -OCOaryl, C<sub>1-6</sub> acyl, C<sub>3-7</sub> membered carbon ring optionally interrupted with 1-4 heteroatoms chosen from O, S, SO, SO<sub>2</sub>, NH and NR<sub>8</sub> where said C<sub>1-6</sub> alkyl, aryl or C<sub>1-6</sub> acyl groups may be independently substituted with 0-3 halogens, hydroxy, N(R)<sub>2</sub>, CO<sub>2</sub>R, C<sub>6-10</sub> aryl, C<sub>5-10</sub> heteroaryl, or C<sub>1-6</sub> alkoxy groups;

20

When two R<sub>13</sub> groups are attached to the same atom or two adjacent atoms they may be taken together to form a 3-7 membered carbon ring optionally interrupted with 1-2 heteroatoms chosen from O, S, SO, SO<sub>2</sub>, NH, and NR<sub>8</sub>;

25

R represents hydrogen, (CH<sub>2</sub>)<sub>p</sub>CN, C<sub>1-6</sub> alkyl, CO<sub>2</sub>C<sub>1-6</sub> alkyl, COCH<sub>2</sub>OH, COCH<sub>2</sub>OCOC<sub>1-6</sub> alkyl, SO<sub>2</sub>C<sub>1-6</sub> alkyl;

30 R<sub>14</sub> represents amino, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, five to six membered heterocycles or phenyl, said phenyl and heterocycles optionally substituted with 1-3 group of halo, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> acylamino, or C<sub>1-6</sub> alkyl, hydroxy and/or amino, said amino and hydroxy optionally protected with an amino or hydroxy protecting group;

R<sub>15</sub> is C<sub>1-6</sub> alkyl or benzyl said benzyl optionally substituted with 1-3 groups of halo, OH, C<sub>1-6</sub> alkoxy, amino, C<sub>1-6</sub> acylamino, or C<sub>1-6</sub> alkyl;

R<sub>16</sub> is hydrogen, C<sub>5-10</sub>heteroaryl, C<sub>6-10</sub>aryl, said heteroaryl and aryl optionally substituted with 1-3 groups of R<sub>7</sub>;

p represents 0-2 and

m, n, and q represents 0-1.

Another aspect of the invention is concerned with the use of the novel antibiotic compositions in the treatment of bacterial infections.

#### DETAILED DESCRIPTION OF THE INVENTION

The invention is described herein in detail using the terms defined below unless otherwise specified.

The compounds of the present invention may have asymmetric centers, chiral axes and chiral planes, and occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers, including optical isomers, being included in the present invention. (See E.L. Eliel and S. H. Wilen Stereochemistry of Carbon Compounds (John Wiley and Sons, New York 1994, in particular pages 1119-1190).

When any variable (e.g. aryl, heterocycle, R<sub>5</sub>, R<sub>6</sub> etc.) occurs more than once, its definition on each occurrence is independent at every other occurrence. Also combinations of substituents/or variables are permissible only if such combinations result in stable compounds.

The term "alkyl" refers to a monovalent alkane (hydrocarbon) derived radical containing from 1 to 15 carbon atoms unless otherwise defined. It may be straight or branched. Preferred alkyl groups include lower alkyls which have from 1 to 6 carbon atoms such as methyl, ethyl, propyl, isopropyl, butyl and t-butyl. When substituted, alkyl groups may be substituted with up to 3 substituent groups, selected from the groups as herein defined, at any available point of attachment. When the alkyl group is said to be substituted with an alkyl group, this is used interchangeably with "branched alkyl group".


Cycloalkyl is a species of alkyl containing from 3 to 15 carbon atoms, without alternating or resonating double bonds between carbon atoms. It may contain from 1 to 4 rings which are fused. Preferred cycloalkyl groups are cyclopropyl,



cyclobutyl, cyclopentyl and cyclohexyl. When substituted, cycloalkyl groups may be substituted with up to 3 substituents which are defined herein by the definition of alkyl.

Alkanoyl refers to a group derived from an aliphatic carboxylic acid of 2 to 4 carbon atoms. Examples are acetyl, propionyl, butyryl and the like.

- 5           The term "alkoxy" refers to those groups of the designated length in either a straight or branched configuration and if two or more carbon atoms in length, they may include a double or a triple bond. Exemplary of such alkoxy groups are methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, tertiary butoxy, pentoxy, isopentoxo, hexoxy, isohexoxy allyloxy, propargyloxy, and the like.




- 10           refers to aryl or heteroaryl, heterocycle, Het, heterocyclyl or heterocyclic as described immediately below.

- Aryl refers to any stable monocyclic or bicyclic carbon ring of up to 7 atoms in each ring, wherein at least one ring is aromatic. Examples of such aryl elements include phenyl, naphthyl, tetrahydronaphthyl, indanyl, indanonyl, biphenyl, tetralinyl, tetralonyl, fluorenyl, phenanthryl, anthryl, acenaphthyl, and the like substituted phenyl and the like. Aryl groups may likewise be substituted as defined. Preferred substituted aryls include phenyl and naphthyl.

- 20           The term heterocycle, heteroaryl, Het, heterocyclyl or heterocyclic, as used herein except where noted, represents a stable 5- to 7-membered mono- or bicyclic or stable 8- to 11-membered bicyclic heterocyclic ring system, any ring of which may be saturated or unsaturated, and which consists of carbon atoms and from one to four heteroatoms selected from the group consisting of N, O and S, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized (in which case it is properly balanced by a counterion), and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom, which results in the creation of a stable structure. The term heterocycle or heterocyclic includes heteroaryl moieties. "Heterocycle" or "heterocyclyl" therefore includes the above mentioned heteroaryls, as well as dihydro and tetrahydro analogs thereof. The heterocycle, heteroaryl, Het or heterocyclic may be substituted with 1-3 groups of R7. Examples of such heterocyclic elements include, but are not limited to the following: piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, 2-oxoazepinyl, azepinyl, pyrrolyl, 4-piperidonyl, pyrrolidinyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazoliny, imidazolidinyl,

- pyridyl, pyrazinyl, pyrimidinyl, pyrimidonyl, pyridinonyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, quinuclidinyl, isothiazolidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, thiadiazolyl, benzopyranyl, benzothiazolyl, benzoxazolyl, furyl, tetrahydrofuryl, 5 tetrahydropyranyl, thiophenyl, imidazopyridinyl, triazolyl, tetrazolyl, triazinyl, thienyl, benzothienyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, naphthpyridinyl, dihydropyrimidinyl, dihydropyrrolyl, dihydroquinolinyl, dihydrotetrazolyl, dihydrotriazolyl, dihydrothienyl, dihydrooxazolyl, dihydrobenzothiophenyl, dihydrofuranlyl, benzothiazolyl, benzothienyl, benzoimidazolyl, benzopyranyl, benzothiofuranyl, carbolinyl, chromanyl, 10 cinnolinyl, benzopyrazolyl, benzodioxolyl and oxadiazolyl. Additional examples of heteroaryls are illustrated by formulas a, b, c and d:



- 15 wherein R<sub>16</sub> and R<sub>17</sub> are independently selected from hydrogen, halogen, C<sub>1-6</sub> alkyl, C<sub>2-4</sub> alkanoyl, C<sub>1-6</sub> alkoxy; and R<sub>18</sub> represents hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-4</sub> alkanoyl, C<sub>1-6</sub> alkoxycarbonyl and carbamoyl.

- The term "alkenyl" refers to a hydrocarbon radical straight, branched or cyclic containing from 2 to 10 carbon atoms and at least one carbon to carbon double bond. 20 Preferred alkenyl groups include ethenyl, propenyl, butenyl and cyclohexenyl.

- The terms "quaternary nitrogen" and "positive charge" refer to tetravalent, positively charged nitrogen atoms (balanced as needed by a counterion known in the art) including, e.g., the positively charged nitrogen in a tetraalkylammonium group (e. g. tetramethylammonium), heteroarylium, (e.g., N-methyl- 25 pyridinium), basic nitrogens which are protonated at physiological pH, and the like. Cationic groups thus encompass positively charged nitrogen-containing groups, as well as basic nitrogens which are protonated at physiologic pH.

The term "heteroatom" means O, S or N, selected on an independent basis.

- The term "prodrug" refers to compounds which are drug precursors which, 30 following administration and absorption, release the drug in vivo via some metabolic process.

Exemplary prodrugs include acyl amides of the amino compounds of this invention such as amides of alkanolic(C<sub>1-6</sub>)acids, amides of aryl acids (e.g., benzoic acid) and alkanedic(C<sub>1-6</sub>)dioic acids.

Halogen and "halo" refer to bromine, chlorine, fluorine and iodine.

- 5                   When a group is termed "substituted", unless otherwise indicated, this means that the group contains from 1 to 3 substituents thereon.

                  When a functional group is termed "protected", this means that the group is in modified form to preclude undesired side reactions at the protected site. Suitable protecting groups for the compounds of the present invention will be recognized from the present application taking into account the level of skill in the art, and with reference to  
10                   standard textbooks, such as Greene, T. W. et al. Protective Groups in Organic Synthesis Wiley, New York (1991). Examples of suitable protecting groups are contained throughout the specification.

                  Examples of suitable hydroxyl and amino protecting groups are:

- 15                   trimethylsilyl, triethylsilyl, o-nitrobenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, t-butylidiphenylsilyl, t-butyldimethylsilyl, benzyloxycarbonyl, t-butyloxycarbonyl, 2,2,2-trichloroethyloxycarbonyl, allyloxycarbonyl and the like. Examples of suitable carboxyl protecting groups are benzhydryl, o-nitrobenzyl, p-nitrobenzyl, 2-naphthylmethyl, allyl, 2-chloroallyl, benzyl, 2,2,2-trichloroethyl, trimethylsilyl, t-butyldimethylsilyl, t-butylidiphenylsilyl, 2-(trimethylsilyl)ethyl, phenacyl, p-methoxybenzyl, acetonyl, p-methoxyphenyl, 4-pyridylmethyl, t-butyl and the like.  
20

- The cyclopropyl containing oxazolidinone compounds of the present invention are useful per se and in their pharmaceutically acceptable salt and ester forms for the treatment of bacterial infections in animal and human subjects. The term  
25                   "pharmaceutically acceptable ester, salt or hydrate," refers to those salts, esters and hydrated forms of the compounds of the present invention which would be apparent to the pharmaceutical chemist. i.e., those which are substantially non-toxic and which may favorably affect the pharmacokinetic properties of said compounds, such as palatability, absorption, distribution, metabolism and excretion. Other factors, more practical in nature, which are also important in the selection, are cost of the raw materials, ease of  
30                   crystallization, yield, stability, solubility, hygroscopicity and flowability of the resulting bulk drug. Conveniently, pharmaceutical compositions may be prepared from the active ingredients in combination with pharmaceutically acceptable carriers. Thus, the present invention is also concerned with pharmaceutical compositions and methods of treating

bacterial infections utilizing as an active ingredient the novel cyclopropyl containing oxazolidinone compounds.

The pharmaceutically acceptable salts referred to above also include acid addition salts. Thus, when the Formula I compounds are basic, salts may be prepared from pharmaceutically acceptable non-toxic acids, including inorganic or organic acids. Included among such acid salts are the following: acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, isethionic, lactate, maleate, mandelic, malic, maleic, methanesulfonate, mucic, 2-naphthalenesulfonate, nicotinate, nitric oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, phosphate, pantothenic, pamoic, sulfate, succinate, tartrate, thiocyanate, tosylate and undecanoate.

When the compound of the present invention is acidic, suitable "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases including inorganic bases and organic bases. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium zinc and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium and sodium salts. Salts derived from pharmaceutically acceptable inorganic non-toxic bases include salts of primary, secondary and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as arginine, betaine caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine tripropylamine, tromethamine and the like.

The pharmaceutically acceptable esters are such as would be readily apparent to a medicinal chemist, and include those which are hydrolyzed under physiological conditions, such as "biolabile esters", pivaloyloxymethyl, acetoxymethyl, phthalidyl, indanyl and methoxymethyl, and others.

Biolabile esters are biologically hydrolyzable, and may be suitable for oral administration, due to good absorption through the stomach or intestinal mucosa, resistance to gastric acid degradation and other factors. Examples of biolabile esters include compounds.

An embodiment of this invention is realized when  $R_1$  represents H,  $NR_5R_6$ , CN, OH,  $C(R_2)OR_{14}$ ,  $NHC(=X_1)N(R_{13})_2$ ,  $C(=NOH)N(R_{13})_2$ ,  $NR_{10}C(=X_1)R_{13}$  or  $CR_7R_8R_9$  and all other variables are as described herein.



Another embodiment of this invention is realized when  
 5 is phenyl, pyridine, pyrimidine, or piperidine and all other variables are as described herein.


Another embodiment of this invention is realized when one of  $R_1$  is  $NR_{10}C(=X_1)R_{13}$  and all other variables are as described herein.

Another embodiment of this invention is realized when one of  $R_1$  is CN and all other variables are as described herein.

10 Another embodiment of this invention is realized when one of  $R_1$  is  $NR_5R_6$  and all other variables are as described herein.

Another embodiment of this invention is realized when  $R_3$  is  $NR(C=X_1)R_{12}$ , C5-10 heteroaryl,  $NH(CH_2)_{0-4}$ aryl,  $NH(CH_2)_{0-4}$ heteroaryl, said aryl and heteroaryl optionally substituted with 1-3 groups of  $R_a$  and all other variables are as  
 15 described herein.

Another embodiment of this invention is realized when  $R_3$  is a C5-10 heteroaryl

represented by  which represents an optionally substituted aromatic heterocyclic group containing 1 to 4 nitrogen atoms and at least one double bond, and which is connected through a bond on any nitrogen. Exemplary groups are 1,2,3-triazole, 1,2,4-triazole, 1,2,5-triazole, tetrazole, pyrazole, and imidazole, any of which may contain 1 to 3 substituents selected from  $R_7$ .  
 20


Still another embodiment of this invention is realized when  $R_5$  and  $R_6$  independently are:

- i) H,
- 25 ii)  $C_{1-6}$  alkyl optionally substituted with 1-3 groups of halogen, CN, OH,  $C_{1-6}$  alkoxy, amino, hydroxyamino, alkoxyamino,  $C_{1-6}$  acyloxy,  $C_{1-6}$  alkylsulfinyl,  $C_{1-6}$  alkylsulfinyl,  $C_{1-6}$  alkylsulfonyl, aminosulfonyl,  $C_{1-6}$  alkylaminosulfonyl,  $C_{1-6}$  dialkylaminosulfonyl, 4-morpholinylsulfonyl, phenyl, pyridine, 5-isoxazolyl, ethenyloxy, or ethynyl, said phenyl and pyridine optionally substituted with 1-3  
 30 halogen, CN, OH,  $CF_3$ ,  $C_{1-6}$  alkyl or  $C_{1-6}$  alkoxy;

- iii) C<sub>1-6</sub> acyl optionally substituted with 1-3 groups of halogen, OH, SH, C<sub>1-6</sub> alkoxy, naphthalenoxy, phenoxy, amino, C<sub>1-6</sub> acylamino, hydroxylamino, alkoxyamino, C<sub>1-6</sub> acyloxy, phenyl, pyridine, C<sub>1-6</sub> alkylcarbonyl, C<sub>1-6</sub> alkylamino, C<sub>1-6</sub> dialkylamino, C<sub>1-6</sub> hydroxyacyloxy, C<sub>1-6</sub> alkylsulfenyl, phthalimido, maleimido, succinimido, said phenoxy, phenyl and pyridine optionally substituted with 1-3 groups of halo, OH, CN, C<sub>1-6</sub> alkoxy, amino, C<sub>1-6</sub> acylamino, CF<sub>3</sub> or C<sub>1-6</sub> alkyl; or
- iv) benzoyl optionally substituted with 1-3 groups of halogen, OH, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkyl, CF<sub>3</sub>, C<sub>1-6</sub> alkanoyl, amino or C<sub>1-6</sub> acylamino and all other variables are as described herein.

Yet another embodiment of this invention is realized when X<sub>1</sub> represents O and all other variables are as described herein.

A preferred embodiment of this invention is realized when the structural formula is II:



Formula II

wherein R<sub>1</sub>, R<sub>4</sub>, R<sub>4a</sub>, Y and R<sub>3</sub> are as described herein.

Preferred compounds of this invention are:

- N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,
- 1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole,

- N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride,
- 5 1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole,  
 1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole hydrochloride,
- N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-acetoxyacetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,
- 10 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-hydroxyacetyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-methanesulfonyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,
- 15 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-methyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3,6-dicyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-cyanomethyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,
- 20 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[(isoxazol-3-yl)oxy]methyloxazolidin-2-one,  
 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[(isoxazol-3-yl)oxy]methyloxazolidin-2-one hydrochloride,
- 25 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[N-(t-butoxycarbonyl)-N-(isoxazol-3-yl)]aminomethyloxazolidin-2-one,  
 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[N-(isoxazol-3-yl)]aminomethyloxazolidin-2-one hydrochloride,
- N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-(5-cyanopyridin-2-yl)-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,
- 30 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-(pyridin-2-yl)-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-acetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,

- N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(pyrimidin-2-yl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(4-pyridylmethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 5 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(N-cyano-1-iminoethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-methoxycarbonyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(N-cyano-S-methylthioiminomethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 10 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(N-cyanocarboxamidyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-(N,N'-t-butoxycarbonylcarboxamidyl)-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 15 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-carboxamidyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-(N-t-Butoxycarbonylamino)acetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-aminoacetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride,  
 20 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-methanesulfonylacetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(dibenzylphosphoryloxy)acetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 25 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(phosphoryloxy)acetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,

or their enantiomer, diastereomer, or pharmaceutically acceptable salt, hydrate or prodrug thereof wherein.

30

Suitable subjects for the administration of the formulation of the present invention include mammals, primates, man, and other animals. *In vitro* antibacterial activity is predictive of *in vivo* activity when the compositions are administered to a mammal infected with a susceptible bacterial organism.



Using standard susceptibility tests, the compositions of the invention are determined to be active against MRSA and enterococcal infections.

The compounds of the invention are formulated in pharmaceutical compositions by combining the compounds with a pharmaceutically acceptable carrier.

5 Examples of such carriers are set forth below.

The compounds may be employed in powder or crystalline form, in liquid solution, or in suspension. They may be administered by a variety of means; those of principal interest include: topically, orally and parenterally by injection (intravenously or intramuscularly).

10 Compositions for injection, a preferred route of delivery, may be prepared in unit dosage form in ampules, or in multidose containers. The injectable compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and may contain various formulating agents. Alternatively, the active ingredient may be in powder (lyophilized or non-lyophilized) form for reconstitution at  
15 the time of delivery with a suitable vehicle, such as sterile water. In injectable compositions, the carrier is typically comprised of sterile water, saline or another injectable liquid, e.g., peanut oil for intramuscular injections. Also, various buffering agents, preservatives and the like can be included.

20 Topical applications may be formulated in carriers such as hydrophobic or hydrophilic bases to form ointments, creams, lotions, in aqueous, oleaginous or alcoholic liquids to form paints or in dry diluents to form powders.

Oral compositions may take such forms as tablets, capsules, oral suspensions and oral solutions. The oral compositions may utilize carriers such as conventional formulating agents, and may include sustained release properties as well as  
25 rapid delivery forms.

The dosage to be administered depends to a large extent upon the condition and size of the subject being treated, the route and frequency of administration, the sensitivity of the pathogen to the particular compound selected, the virulence of the infection and other factors. Such matters, however, are left to the routine discretion of  
30 the physician according to principles of treatment well known in the antibacterial arts. Another factor influencing the precise dosage regimen, apart from the nature of the infection and peculiar identity of the individual being treated, is the molecular weight of the compound.

The novel antibiotic compositions of this invention for human delivery  
35 per unit dosage, whether liquid or solid, comprise from about 0.01% to as high as about

99% of the cyclopropyl containing oxazolidinone compounds discussed herein, the preferred range being from about 10-60% and from about 1% to about 99.99% of one or more of other antibiotics such as those discussed herein, preferably from about 40% to about 90%. The composition will generally contain from about 125 mg to about 3.0 g of the cyclopropyl containing oxazolidinone compounds discussed herein; however, in  
5      general, it is preferable to employ dosage amounts in the range of from about 250 mg to 1000 mg and from about 200mg to about 5 g of the other antibiotics discussed herein; preferably from about 250 mg to about 1000 mg. In parenteral administration, the unit dosage will typically include the pure compound in sterile water solution or in the form  
10     of a soluble powder intended for solution, which can be adjusted to neutral pH and isotonic.

The invention described herein also includes a method of treating a bacterial infection in a mammal in need of such treatment comprising administering to said mammal the claimed composition in an amount effective to treat said infection.

15       Oxazolidinones have been known at times to cause side effects such as sideroblastic anemia, peripheral sensory neuropathy, optic neuropathy, seizures, thrombocytopenia, cheilosis, seborrheic dermatitis, hypo-regenerative anemia, megaloblastic anemia or normocytic anemia. The compounds of the invention may be combined with an effective amount of one or more vitamins to prevent or reduce the  
20     occurrence of oxazolidinone-associated side effects in patients. The vitamins that can be combined are vitamin B2, vitamin B6, vitamin B12 and folic acid. The vitamins may be administered with the oxazolidinones as separate compositions or the vitamins and oxazolidinones may be present in the same composition.

Thus another aspect of this invention is a method of treating or  
25     preventing an oxazolidinone-associated side effect by administering an effective amount of the oxazolidinone of structural formula I and an effective amount of one or more of vitamin B2, vitamin B6, vitamin B12 and folic acid to a patient in need thereof.

A further aspect of this invention relates to a method of treating or preventing oxazolidinone-associated normocytic anemia or peripheral sensory  
30     neuropathy by administering an effective amount of vitamin B2 to a patient in need thereof.

Yet another aspect of this invention relates to a method of treating or preventing oxazolidinone-associated sideroblastic anemia, peripheral sensory neuropathy, optic neuropathy, seizures, thrombocytopenia, cheilosis, and seborrheic

dermatitis by administering an effective amount of vitamin B6 to a patient in need thereof.

Still another aspect of this invention relates to a method of treating or preventing oxazolidinone-associated hypo-regenerative anemia, megaloblastic anemia by  
 5 administering an effective amount of vitamin B12 and folic acid to a patient in need thereof.

Still another aspect of this invention relates to a method of treating or preventing bacterial infection by administering an effective amount of a compound of formula I and an effective amount of one or more of the group selected from the group  
 10 consisting of vitamin B2, vitamin B6, vitamin B12 and folic acid to a patient in need thereof.


The preferred methods of administration of the claimed compositions include oral and parenteral, e.g., i.v. infusion, i.v. bolus and i.m. injection formulated so that a unit dosage comprises a therapeutically effective amount of each active  
 15 component or some submultiple thereof.

For adults, about 5-50 mg/kg of body weight, preferably about 250 mg to about 1000 mg per person of the cyclopropyl containing oxazolidinone antibacterial compound and about 250 mg, to about 1000 mg per person of the other antibiotic(s) given one to four times daily is preferred. More specifically, for mild infections a dose  
 20 of about 250 mg two or three times daily of the cyclopropyl containing oxazolidinone antibacterial compound and about 250 mg two or three times daily of the other antibiotic is recommended. For moderate infections against highly susceptible gram positive organisms a dose of about 500 mg each of the cyclopropyl containing oxazolidinone and the other antibiotics, three or four times daily is recommended. For severe, life-  
 25 threatening infections against organisms at the upper limits of sensitivity to the antibiotic, a dose of about 500-2000 mg each of the cyclopropyl-containing oxazolidinone compound and the other antibiotics, three to four times daily may be recommended.

For children, a dose of about 5-25 mg/kg of body weight given 2, 3, or 4  
 30 times per day is preferred; a dose of 10 mg/kg is typically recommended.

The invention is further described in connection with the following non-limiting examples.

#### EXAMPLE 1



N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

Step 1.

- 5        5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]-5-hydroxymethyloxazolidin-2-one.

To a solution of 1-benzyloxycarbonylamino-4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]benzene (1.26 g) in dry tetrahydrofuran (25 mL) was added a solution of n-butyllithium in hexane (1.6 M, 2.51 mL) at -78 °C, and the mixture was stirred at the same temperature for 30 min. (R)-Glycidyl butyrate (0.58 mL) was added to the mixture at -78 °C and the mixture was stirred at room temperature for 2 hours. After quenching the reaction with the addition of methanol (2.5 mL), the mixture was stirred at room temperature for 30 minutes. After dilution of the mixture with aqueous ammonium chloride solution, the mixture was extracted with ethyl acetate. The organic extracts were washed with brine, dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Flash chromatography (silica, ethyl acetate) of the residue gave 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]-5-hydroxymethyloxazolidin-2-one (995 mg).

MS (EI<sup>+</sup>) *m/z*: 298 (M<sup>+</sup>).

- 20    HRMS (EI<sup>+</sup>) for C<sub>17</sub>H<sub>18</sub>N<sub>2</sub>O<sub>3</sub> (M<sup>+</sup>): calcd, 298.1317; found, 298.1310.

Step 2.

5(R)-Azidomethyl-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]oxazolidin-2-one.

- 25    To a solution of 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]-5-hydroxymethyloxazolidin-2-one (298 mg) in dichloromethane (10 mL) was added triethylamine (0.28 mL) and methanesulfonyl chloride (0.12 mL) at 0 °C, the mixture was stirred at the same temperature for 15 minutes. After dilution of the mixture with 1 N hydrochloric acid, the mixture was extracted with ethyl acetate. The organic extracts were washed with water, aqueous sodium hydrogencarbonate solution and brine, dried over


anhydrous sodium sulfate, filtered, and then concentrated in vacuo. A suspension of the residue and sodium azide (199 mg) in N,N-dimethylformamide (10 mL) was stirred at 70 °C for 4 hours and concentrated in vacuo. After dilution of the residue with water, the mixture extracted with ethyl acetate. The organic extracts were washed with water and brine, dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Flash chromatography (silica, ethyl acetate) of the residue gave 5(R)-azidomethyl-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]oxazolidin-2-one (304 mg). MS (EI<sup>+</sup>) *m/z*: 323 (M<sup>+</sup>). HRMS (EI<sup>+</sup>) for C<sub>17</sub>H<sub>17</sub>N<sub>3</sub>O<sub>2</sub> (M<sup>+</sup>): calcd, 323.1382; found, 323.1363.

10 Step 3.

N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

A suspension of 5(R)-azidomethyl-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]oxazolidin-2-one (300 mg) and Lindlar catalyst (5% palladium on CaCO<sub>3</sub> partially poisoned with lead, 150 mg) in tetrahydrofuran (2 mL) and methanol (10 mL) was hydrogenated at 1 atm for 70 minutes at room temperature. After filtration of the catalyst, the filtrate was concentrated in vacuo to give 5(R)-aminomethyl-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]oxazolidin-2-one (276 mg). This compound was used without further purification. To a solution of the crude 5(R)-aminomethyl-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]oxazolidin-2-one (276 mg) in tetrahydrofuran (10 mL) was added triethylamine (194  $\mu$ L) and acetic anhydride (108  $\mu$ L) at 0 °C, and the mixture was stirred at room temperature for 30 minutes. After quenching the reaction by the addition of 1 N hydrochloric acid, the mixture was extracted with ethyl acetate. The organic extracts were washed with water, aqueous sodium hydrogencarbonate solution and brine, dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Flash chromatography (silica, ethyl acetate : methanol = 15:1) of the residue gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (276 mg). MS (EI<sup>+</sup>) *m/z*: 339 (M<sup>+</sup>). HRMS (EI<sup>+</sup>) for C<sub>19</sub>H<sub>21</sub>N<sub>3</sub>O<sub>3</sub> (M<sup>+</sup>): calcd, 339.1583; found, 339.1606.

EXAMPLE 2




1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole.

The mixture of 5(R)-azidomethyl-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]oxazolidin-2-one (417 mg) and 2,5-norbornadiene (0.70 mL) in dioxane (13 mL) was heated under reflux for 4 hours, and then concentrated in vacuo. Flash chromatography (silica, ethyl acetate : methanol = 20:1) of the residue gave 1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole (345 mg).

MS (EI<sup>+</sup>) *m/z*: 349 (M<sup>+</sup>).

HRMS (EI<sup>+</sup>) for C<sub>19</sub>H<sub>19</sub>FN<sub>5</sub>O<sub>2</sub> (M<sup>+</sup>): calcd, 349.1539; found, 349.1526.

### EXAMPLE 3



N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-Butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

#### Step 1.

5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-Butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-hydroxymethyloxazolidin-2-one.

The title compound 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-hydroxymethyloxazolidin-2-one (102 mg) was prepared from 4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]-

1-benzyloxycarbonylaminobenzene (123 mg) in the same manner as described for EXAMPLE 1.

MS (EI<sup>+</sup>) *m/z*: 399 (M<sup>+</sup>).

HRMS (EI<sup>+</sup>) for C<sub>21</sub>H<sub>23</sub>N<sub>3</sub>O<sub>3</sub> (M<sup>+</sup>): calcd, 399.1794; found, 399.1801.

5 Step 2.

N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-Butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.


The title compound N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (89.9 mg) was prepared from 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-hydroxymethyloxazolidin-2-one (98.4 mg) in the same manner as described for EXAMPLE 1.

MS (EI<sup>+</sup>) *m/z*: 440 (M<sup>+</sup>).

HRMS (EI<sup>+</sup>) for C<sub>23</sub>H<sub>28</sub>N<sub>4</sub>O<sub>3</sub> (M<sup>+</sup>): calcd, 440.2060; found, 440.2076.

15

EXAMPLE 4




N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide Hydrochloride.

20 To a solution of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (378 mg) in tetrahydrofuran (5 mL) was added a solution of hydrogen chloride in ethanol (10 M, 15 mL) at 0 °C, the mixture was stirred at room temperature for 3 hours and concentrated in vacuo. Treatment with ethanol of the residue gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (275 mg).

MS (EI<sup>+</sup>) *m/z*: 340 (M<sup>+</sup>) (as free base).

HRMS (EI<sup>+</sup>) for C<sub>18</sub>H<sub>20</sub>N<sub>4</sub>O<sub>3</sub> (M<sup>+</sup>): calcd, 340.1535; found, 340.1553.


**EXAMPLE 5**

1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-Butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole.

- 5 The title compound 1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole (358 mg) was prepared from 5(R)-azidomethyl-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]oxazolidin-2-one (425 mg) in the same manner as described for EXAMPLE 2.

10 MS (FAB<sup>+</sup>) *m/z*: 451 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>27</sub>N<sub>6</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 451.2094; found, 451.2098.

**EXAMPLE 6**

- 15 1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole Hydrochloride.


1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole hydrochloride (267 mg) was prepared from 1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole (358 mg) in the same manner as described for EXAMPLE 4.



MS (EI<sup>+</sup>) *m/z*: 350 (M<sup>+</sup>) (as free base).

HRMS (EI<sup>+</sup>) for C<sub>18</sub>H<sub>18</sub>N<sub>6</sub>O<sub>2</sub> (M<sup>+</sup>): calcd, 350.1491; found, 350.1464.

#### EXAMPLE 7



5

N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-Acetoxyacetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.


To a suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (415 mg) in  
 10 dichloromethane (11 mL) was added triethylamine (0.46 mL) and acetoxyacetyl chloride (0.15 mL) at 0 °C, the mixture was stirred at the same temperature for 45 minutes. After dilution of the mixture with water, the mixture was extracted with dichloromethane. The organic extracts were dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Flash chromatography (silica, dichloromethane : methanol = 10:1) of the residue gave  
 15 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-acetoxyacetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (378 mg).

MS (FAB<sup>+</sup>) *m/z*: 441 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>25</sub>N<sub>4</sub>O<sub>6</sub> (MH<sup>+</sup>): calcd, 441.1774; found, 441.1764.

20


#### EXAMPLE 8



N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyano-3-hydroxyacetyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.


- To a suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-acetoxyacetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (225 mg) in methanol (5 mL) and tetrahydrofuran (1 mL) was added potassium carbonate (141 mg) at room temperature, the mixture was stirred at the same temperature for 90 minutes and concentrated in vacuo. Flash chromatography (silica, dichloromethane : methanol = 20:1) of the residue gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-hydroxyacetyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (138 mg).
- MS (FAB<sup>+</sup>) *m/z*: 399 (MH<sup>+</sup>).
- HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>23</sub>N<sub>4</sub>O<sub>5</sub> (MH<sup>+</sup>): calcd, 399.1668; found, 399.1681.

#### EXAMPLE 9



- N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyano-3-methanesulfonyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.
- The title compound N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-methanesulfonyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (219 mg) was prepared from N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (226 mg) and methanesulfonyl chloride (70  $\mu$ L) in the same manner as described for EXAMPLE 7.
- MS (FAB<sup>+</sup>) *m/z*: 419 (MH<sup>+</sup>).
- HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>23</sub>N<sub>4</sub>O<sub>5</sub>S (MH<sup>+</sup>): calcd, 419.1389; found, 419.1386.


#### EXAMPLE 10



N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-Cyano-3-methyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

- To a suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (188 mg) in tetrahydrofuran (5 mL) was added acetic acid (57  $\mu$ L), 35 % formaldehyde (396  $\mu$ L), and sodium triacetoxyborohydride (223 mg) at room temperature, the mixture was stirred at the same temperature for 2 hours. After quenching the reaction by addition of aqueous sodium hydrogencarbonate solution, the mixture was extracted with dichloromethane-methanol (5:1).
- The organic extracts were dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Flash chromatography (silica, dichloromethane : methanol = 10:1) of the residue gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-methyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (104 mg).
- MS (FAB<sup>+</sup>) *m/z*: 355 (MH<sup>+</sup>).
- HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>23</sub>N<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 355.1770; found, 355.1775.

#### EXAMPLE 11



- N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3,6-Dicyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.


A suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (245 mg) and sodium acetate (373 mg) in methanol (22 mL) was stirred at room temperature for 20 minutes. To

the resulting suspension was added a solution of cyanogens bromide in dichloromethane (5 M, 0.26 mL) at 0°C, the mixture was stirred at the same temperature for 40 minutes, and concentrated in vacuo. Flash chromatography (silica, dichloromethane : methanol = 10:1) of the residue gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3,6-dicyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (207 mg).

MS (FAB<sup>+</sup>) *m/z*: 366 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>20</sub>N<sub>3</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 366.1566; found, 366.1575.

#### EXAMPLE 12




N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyano-3-cyanomethyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

A suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-cyanomethyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (245 mg), sodium hydrogencarbonate (273 mg) in N,N-dimethylformamide (6.5 mL) was stirred at room temperature for 10 minutes. To the resulting suspension was added bromoacetonitrile (70  $\mu$ L) at room temperature, the mixture was stirred at the same temperature for 6 hours, and concentrated in vacuo. Flash chromatography (silica, dichloromethane : methanol = 10:1) of the residue gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-cyanomethyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (219 mg).

MS (FAB<sup>+</sup>) *m/z*: 380 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>22</sub>N<sub>3</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 380.1723; found, 380.1728.

#### EXAMPLE 13




5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-Butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[(isoxazol-3-yl)oxy]methyloxazolidin-2-one.

To a suspension of 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-hydroxymethyloxazolidin-2-one (10.0 mg), 3-hydroxyisoxazole (4.3 mg) and triphenylphosphine (13.5 mg) in tetrahydrofuran (0.25 mL) was added diisopropyl azodicarboxylate (9.8  $\mu$ L), the mixture was stirred at room temperature for 3 hours, and concentrated in vacuo. Flash chromatography (silica, hexane : ethyl acetate = 1:5) of the residue gave 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[(isoxazol-3-yl)oxy]methyloxazolidin-2-one (11.7 mg).

MS (FAB<sup>+</sup>)  $m/z$ : 467 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>27</sub>N<sub>4</sub>O<sub>6</sub> (MH<sup>+</sup>): calcd, 467.1931; found, 467.1903.

#### 15 EXAMPLE 14



5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[(isoxazolyl-3-yl)oxy]methyloxazolidin-2-one Hydrochloride.

The title compound 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[(isoxazolyl-3-yl)oxy]methyloxazolidin-2-one hydrochloride (199 mg) was prepared from 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-


azabicyclo[3.1.0]hexan-6-yl]]phenyl]-5-[(isoxazolyl-3-yl)oxy]methyloxazolidin-2-one (248 mg) in the same manner as described for EXAMPLE 4.

MS (EI<sup>+</sup>) *m/z*: 366 (M<sup>+</sup>) (as free base).

HRMS (EI<sup>+</sup>) for C<sub>19</sub>H<sub>18</sub>N<sub>4</sub>O<sub>4</sub> (M<sup>+</sup>): calcd, 366.1328; found, 366.1330.

5

#### EXAMPLE 15




5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-Butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[N-(t-butoxycarbonyl)-N-(isoxazol-3-yl)]aminomethyloxazolidin-2-one.

10 To a suspension of 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-hydroxymethyloxazolidin-2-one (10.0 mg), 3-N-(t-butoxycarbonyl)aminoisoxazole (9.2 mg), and tetramethylazodicarboxamide (8.6 mg) in benzene (0.25 mL) was added tributylphosphine (12.5  $\mu$ L), and the mixture was stirred at room temperature for 90 minutes. After dilution of the mixture with ethyl acetate, the insoluble materials were filtered off, and the filtrate was concentrated in vacuo. Flash chromatography (silica, hexane : ethyl acetate = 3:5) of the mixture gave 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[N-(t-butoxycarbonyl)-N-(isoxazol-3-yl)]aminomethyloxazolidin-2-one (14.1 mg).

15 MS (FAB<sup>+</sup>) *m/z*: 566 (MH<sup>+</sup>).

20 HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>36</sub>N<sub>5</sub>O<sub>7</sub> (MH<sup>+</sup>): calcd, 566.2615; found, 566.2609.

#### EXAMPLE 16




5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[N-(isoxazol-3-yl)]aminomethyloxazolidin-2-one Hydrochloride.

The title compound 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[N-(isoxazol-3-yl)]aminomethyloxazolidin-2-one hydrochloride (207 mg) was prepared from (R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[N-(t-butoxycarbonyl)-N-(isoxazol-3-yl)]aminomethyloxazolidin-2-one (292 mg) in the same manner as described for EXAMPLE 4.

MS (EI<sup>+</sup>) *m/z*: 365 (M<sup>+</sup>) (as free base).

10 HRMS (EI<sup>+</sup>) for C<sub>19</sub>H<sub>19</sub>N<sub>5</sub>O<sub>3</sub> (M<sup>+</sup>): calcd, 365.1488; found, 365.1478.

#### EXAMPLE 17



15 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyano-3-(5-cyanopyridin-2-yl)-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

To a suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (226 mg) in dimethyl sulfoxide (6 mL) was added diisopropylethylamine (1.05 mL), the mixture was stirred at room temperature for 5 minutes. To the resulting mixture was added 2-chloro-5-cyanopyridine (166 mg), the mixture was stirred at 40°C for overnight, and stirred at 60°C for 10 hours. After dilution of the mixture with ethyl acetate and water, the mixture was


extracted with ethyl acetate. The organic extracts were washed with aqueous sodium hydrogencarbonate solution and brine, dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Flash chromatography (silica, dichloromethane : methanol = 10:1) of the residue gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-(5-cyanopyridin-2-yl)-3-

5 azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (218 mg).

MS (FAB<sup>+</sup>) *m/z*: 443 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>22</sub>N<sub>6</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 443.1832; found, 443.1841.

#### EXAMPLE 18



10

N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyano-3-(pyridin-2-yl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

To a suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (264 mg) in 2-pyridyl trifluoromethanesulfonate (5.49 mL) was added diisopropylethylamine (1.22 mL), the mixture was stirred at room temperature for 5 minutes, and stirred at 90°C for 30.5 hours. Flash chromatography (silica, ethyl acetate: methanol = 5:1) of the mixture gave N-[5(S)-3-

15


[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-(pyridin-2-yl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (215 mg).

20 MS (FAB<sup>+</sup>) *m/z*: 418 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>24</sub>N<sub>3</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 418.1879; found, 418.1885.

#### EXAMPLE 19






N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-Acetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

To a suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (245 mg) in tetrahydrofuran (6.5 mL) was added saturated sodium hydrogencarbonate solution (6.5 mL), the mixture was stirred at 0°C for 5 minutes. To the resulting mixture was added acetic anhydride (70  $\mu$ L), the mixture was and stirred at 0°C for 20 minutes. After an aqueous layer of the mixture was extracted with dichloromethane-methanol (10:1), the combined organic  
 10 extracts were dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Treatment of the residue with ethyl acetate gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-acetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (215 mg).

MS (FAB<sup>+</sup>) *m/z*: 383 (MH<sup>+</sup>).

15 HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>23</sub>N<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 383.1719; found, 383.1732.

#### EXAMPLE 20




20 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-Cyano-3-(pyrimidin-2-yl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

The title compound N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-(pyrimidin-2-yl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (198 mg) was prepared from N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (283 mg) and 2-chloropyrimidine (181 mg) in the same manner as described for EXAMPLE 17.

MS (FAB<sup>+</sup>) *m/z*: 419 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>23</sub>N<sub>6</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 419.1832; found, 419.1832.

#### EXAMPLE 21




10

N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyano-3-(4-pyridylmethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

To a suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (188 mg) in dichloromethane (33 mL) was added triethylamine (209  $\mu$ L), the mixture was stirred at room temperature for 5 minutes. To the resulting mixture was added pyridine 4-carboxamide (98  $\mu$ L), acetic acid (115  $\mu$ L), and sodium triacetoxyborohydride (223 mg) at room temperature, the mixture was stirred at room temperature for 7 hours. After quenching the reaction by addition of 1N sodium hydroxide solution, the mixture was extracted with dichloromethane. The organic extracts were dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Flash chromatography (silica, dichloromethane : methanol = 10:1) of the residue gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-(4-pyridylmethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (181 mg). MS (FAB<sup>+</sup>) *m/z*: 432 (MH<sup>+</sup>). HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>26</sub>N<sub>5</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 432.2036; found, 432.2041.

25


#### EXAMPLE 22



N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(N-cyano-1-iminoethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

- To a suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (207 mg) in methanol (5.5 mL) was added diisopropylethylamine (192  $\mu$ L), the mixture was stirred at room temperature for 20 minutes. To the resulting mixture was added methyl N-cyanoacetimidate (108 mg), the mixture was stirred at room temperature for 2 days. The resulting precipitates were collected by filtration to give N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(N-cyano-1-iminoethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (186 mg).  
 MS (FAB<sup>+</sup>) *m/z*: 407 (MH<sup>+</sup>).  
 HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>23</sub>N<sub>6</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 407.1832; found, 407.1869.

#### EXAMPLE 23




- N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-methoxycarbonyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.  
 To a suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (188 mg) in acetonitrile (5 mL) was added diisopropylethylamine (261  $\mu$ L), the mixture was stirred at room temperature for 10 minutes. To the resulting mixture was added methyl chloroformate (61  $\mu$ L) at 0°C, the mixture was stirred at room temperature for 25 minutes, and concentrated in vacuo. After

dilution of the residue with 1N hydrochloric acid, the mixture was extracted with dichloromethane. The organic extracts were dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Treatment of the residue with ethyl acetate gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-methoxycarbonyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (161 mg).

MS (FAB<sup>+</sup>) *m/z*: 399 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>23</sub>N<sub>4</sub>O<sub>5</sub> (MH<sup>+</sup>): calcd, 399.1668; found, 399.1671.

#### EXAMPLE 24




N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-Cyano-3-(N-cyano-S-methylthioiminomethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

The title compound N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(N-cyano-S-methylthioiminomethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (8.3 mg) was prepared from N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (9.4 mg) and dimethyl N-cyanodithioiminocarbonate (4.9 mg) in the same manner as described for EXAMPLE 11.

MS (FAB<sup>+</sup>) *m/z*: 439 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>23</sub>N<sub>6</sub>O<sub>3</sub>S (MH<sup>+</sup>): calcd, 439.1552; found, 439.1553.

#### EXAMPLE 25




N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-Cyano-3-(N-cyanocarboxamidyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

To a solution of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(N-cyano-S-  
 5 methylthioiminomethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (260 mg) in dimethylformamide (8 mL) was added a solution of ammonia in methanol (4.1 M, 8 mL) at 0°C, the mixture was allowed stand at room temperature for 3 days, and concentrated in vacuo. Flash chromatography (silica, dichloromethane : methanol = 20:3) of the residue gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-  
 10 3-(N-cyanocarboxamidyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (111 mg).

MS (FAB<sup>+</sup>) *m/z*: 408 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>22</sub>N<sub>3</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 408.1784; found, 408.1792.

#### 15 EXAMPLE 26



N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-(N,N'-t-Butoxycarbonylcarboxamidyl)-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.


The title compound N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-(N,N'-t-  
 20 butoxycarbonylcarboxamidyl)-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (35 mg) was prepared from N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-

(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (19 mg) and N,N'-di(t-butoxycarbonyl)-1H-pyrazole-1-carboxamidine (23 mg) in the same manner as described for EXAMPLE 23.

MS (FAB<sup>+</sup>) *m/z*: 583 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>39</sub>N<sub>6</sub>O<sub>7</sub> (MH<sup>+</sup>): calcd, 583.2880; found, 583.2880.

#### EXAMPLE 27




10 N-[5(S)-3-[4-[(1α,5α,6β)-[3-Carboxamidyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide Hydrochloride.

To a suspension of N-[5(S)-3-[4-[(1α,5α,6β)-[3-(N,N'-t-butoxycarbonylcarboxamidyl)-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (221 mg) in tetrahydrofuran (1.9 mL) was added a solution of hydrogen chloride in dioxane (4.8 M, 5.7 mL) at 0°C, the mixture was stirred at  
 15 room temperature for 3.7 hours, and concentrated in vacuo. After dilution of the residue with water, the mixture was washed with ethyl acetate. The resulting aqueous layer was lyophilized to give N-[5(S)-3-[4-[(1α,5α,6β)-[3-carboxamidyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (157 mg).

20 MS (FAB<sup>+</sup>) *m/z*: 383 (MH<sup>+</sup>)(as free base).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>23</sub>N<sub>6</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 383.1832; found, 383.1879.


#### EXAMPLE 28



N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-(N-t-Butoxycarbonylamino)acetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

- To a suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (19 mg), N-t-butoxycarbonylglycine (9.6 mg), and 1-hydroxybenzotriazole (8.4 mg) in dimethylformamide (2 mL) was added triethylamine (17  $\mu$ L), the mixture was stirred at room temperature for 5 minutes. The resulting mixture was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (11 mg) at 0°C, the mixture was stirred at room temperature for 5.5 hours, and concentrated in vacuo. After dilution of the residue with 1N hydrochloric acid, the mixture was extracted with ethyl acetate. The organic extracts were washed with aqueous sodium hydrogencarbonate solution, dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Flash chromatography (silica, ethyl acetate : methanol = 5:1) of the residue gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-(N-t-Butoxycarbonylamino)acetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (23 mg). MS (FAB<sup>+</sup>)  $m/z$ : 498 (MH<sup>+</sup>). HRMS (FAB<sup>+</sup>) for C<sub>25</sub>H<sub>32</sub>N<sub>5</sub>O<sub>6</sub> (MH<sup>+</sup>): calcd, 498.2353; found, 498.2339.

#### EXAMPLE 29



20

N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-Aminoacetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide Hydrochloride.


The title compound N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-aminoacetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (225 mg) was prepared from N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-(N-t-butoxycarbonylamino)acetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (299 mg) in the same manner as described for EXAMPLE 27.

MS (FAB<sup>+</sup>) *m/z*: 398 (MH<sup>+</sup>)(as free base).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>24</sub>N<sub>3</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 398.1828; found, 398.1826.

10

#### EXAMPLE 30



N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-Cyano-3-methanesulfonylacetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.


The title compound N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-methanesulfonylacetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (193 mg) was prepared from N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride (188 mg) and methanesulfonylacetic acid (78 mg) in the same manner as described for EXAMPLE 28.

MS (FAB<sup>+</sup>) *m/z*: 461 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>23</sub>N<sub>4</sub>O<sub>6</sub>S (MH<sup>+</sup>): calcd, 461.1495; found, 461.1513.

#### EXAMPLE 31






N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-Cyano-3-(dibenzylphosphoryloxy)acetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

To a suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-hydroxyacetyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (10 mg), triphenylphosphine (14 mg), and dibenzyl phosphate (14 mg) in tetrahydrofuran (1 mL) was added an solution of diisopropyl azodicarboxylate in toluene (40wt%, 27  $\mu$ L), the mixture was stirred at room temperature for overnight, and stirred at 60°C for 11 hours. The mixture was concentrated in vacuo. Flash chromatography (silica, dichloromethane : methanol = 20:3) of the residue gave N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(dibenzylphosphoryloxy)acetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (9.5 mg).

MS (FAB<sup>+</sup>)  $m/z$ : 659 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>34</sub>H<sub>36</sub>N<sub>4</sub>O<sub>8</sub>P (MH<sup>+</sup>): calcd, 659.2271; found, 659.2256.

#### EXAMPLE 32



N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-Cyano-3-(phosphoryloxy)acetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide.

- A suspension of N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(dibenzylphosphoryloxy)acetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (163 mg) and palladium on charcoal (7.5%, 24 mg) in methanol (6 mL) was stirred at room temperature for 4 hours under hydrogen atmosphere. After insoluble materials were filtered off, the filtrate was concentrated in vacuo. A solution of the residue in water (1.5 mL) was washed with ethyl acetate, the resulting aqueous solution was lyophilized to give N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(phosphoryloxy)acetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide (111 mg). MS (FAB<sup>+</sup>) *m/z*: 479 (MH<sup>+</sup>).
- 10 HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>24</sub>N<sub>4</sub>O<sub>8</sub>P (MH<sup>+</sup>): calcd, 479.1332; found, 479.1344.

#### REFERENCE EXAMPLE 1

4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyanobicyclo[3.1.0]hexan-6-yl)]-1-benzyloxycarbonylaminobenzene.

##### 15 Step 1.

(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyanobicyclo[3.1.0]hexan-6-yl)benzene.

- To a solution of lithium diisopropylamide (prepared from diisopropylamine (3.88 mL) and n-butyllithium (1.6 M solution in hexane, 17.4 mL)) in tetrahydrofuran (37 mL) was added phenylacetonitrile (3.18 mL) at -50 °C, the mixture was stirred at 20 0 °C for 3 hours. To the mixture was added a solution of cyclopenten-1-yl phenyl sulfone (5.49 g) in tetrahydrofuran (26 mL) at 5 °C, the mixture was stirred at the same temperature for 40 minutes, and stirred at room temperature for 18 hours. The mixture was stirred at 60 °C for 3 hours. After dilution of the mixture with aqueous ammonium chloride solution, the mixture was extracted with ethyl acetate. The 25 organic extracts were washed with brine, dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Flash chromatography (silica, hexane : ethyl acetate = 10:1) of the residue gave (1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)benzene (4.44 g). MS (EI<sup>+</sup>) *m/z*: 183 (M<sup>+</sup>).

- 30 HRMS (EI<sup>+</sup>) for C<sub>13</sub>H<sub>13</sub>N (M<sup>+</sup>): calcd, 183.1048; found, 183.1072.

##### Step 2.

4-(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)-1-nitrobenzene.

To a solution of (1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)benzene (916 mg) in chloroform (5 mL) was added concentrated sulfuric acid (1.93 mL) and nitric

acid (fuming, 0.28 mL) at  $-30\text{ }^{\circ}\text{C}$ , the mixture was stirred at the same temperature for 1 minute. The mixture was poured into ice water, extracted with chloroform. The organic extracts were washed with aqueous sodium hydrogencarbonate solution and brine, dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo.

- 5 Flash chromatography (silica, hexane : ethyl acetate = 10:3) of the residue gave 4-(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)-1-nitrobenzene (875 mg).

MS ( $\text{EI}^+$ )  $m/z$ : 228 ( $\text{M}^+$ ).

HRMS ( $\text{EI}^+$ ) for  $\text{C}_{13}\text{H}_{12}\text{N}_2\text{O}_2$  ( $\text{M}^+$ ): calcd, 228.0899; found, 228.0889.

### Step 3.

- 10 1-Benzyloxycarbonylamino-4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]benzene.

A suspension of 4-(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)-1-nitrobenzene (875 mg) and palladium catalyst (10 % on chacoal, 87 mg) in tetrahydrofuran (19 mL) was hydrogenated at 1 atm for 3 hours at room temperature.

- 15 After filtration of the catalyst, the filtrate was concentrated in vacuo to give 1-amino-4-(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)benzene. To a solution of crude 1-amino-4-(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)benzene thus obtained in acetone (12 mL) was added sodium hydrogencarbonate (644 mg), water (6 mL) and benzyl chloroformate (0.69 mL) at  $0\text{ }^{\circ}\text{C}$ , the mixture was stirred at the same  
20 temperature for 5 minutes. After dilution of the mixture by addition of ice water, the mixture was extracted with ethyl acetate. The organic extracts were washed with brine, dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Flash chromatography (silica, hexane: ethyl acetate = 5:2) of the residue gave 1-benzyloxycarbonylamino-4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]benzene  
25 (1.27 g).

MS ( $\text{EI}^+$ )  $m/z$ : 332 ( $\text{M}^+$ ).

HRMS ( $\text{EI}^+$ ) for  $\text{C}_{21}\text{H}_{20}\text{N}_2\text{O}_2$  ( $\text{M}^+$ ): calcd, 332.1525; found, 332.1543.

### REFERENCE EXAMPLE 2

- 30 1-t-Butoxycarbonyl-3-pyrrolin-3-yl phenyl sulfone.

To a suspension of N-chlorosuccinimide (781 mg) in dichloromethane (6 mL) was added benzenethiol (0.60 mL) at room temperature, the mixture was stirred at the same temperature for 30 minutes. To the resulting mixture was added a solution of 1-t-butoxycarbonyl-3-pyrroline (1.00 g) in dichloromethane (1 mL) at  $-60\text{ }^{\circ}\text{C}$ , the  
35 mixture was stirred at room temperature for 1 hour. The insoluble materials were

filtered off, the filtrate was concentrated in vacuo. To a solution of the residue in dichloromethane (29 mL) was added m-chloroperoxybenzoic acid (3.65 g) at 0 °C, the mixture was stirred at room temperature for 1 hour. To the resulting mixture was added sodium carbonate (2.19 g), the mixture was stirred at room temperature for 5 minutes. The insoluble materials were filtered off, the filtrate was diluted with ether. The filtrate was washed with 10 % sodium bisulfate solution, 10 % sodium carbonate solution and brine, dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. To a solution of the residue in dichloromethane (11 mL) was added 1,8-diazabicyclo[5.4.0]undec-7-ene (0.92 mL) at -40 °C, the mixture was stirred at room temperature for 5 minutes. The mixture was poured into 1 N hydrochloric acid and extracted with ether. The organic extracts were washed with water, aqueous sodium hydrogencarbonate solution and brine, dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. Flash chromatography (silica, hexane: ethyl acetate = 5:2) of the residue gave 1-t-butoxycarbonyl-3-pyrrolin-3-yl phenyl sulfone (1.16 g).

MS (EI<sup>+</sup>) *m/z*: 309 (M<sup>+</sup>).  
 HRMS (EI<sup>+</sup>) for C<sub>15</sub>H<sub>19</sub>NO<sub>4</sub>S (M<sup>+</sup>): calcd, 309.1035; found, 309.1042.

### REFERENCE EXAMPLE 3

1-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-Butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]-4-nitrobenzene.

#### Step 1.

(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-Butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)benzene.

The title compound (1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)benzene (3.83 g) was prepared from phenylacetonitrile (1.65 mL) and 1-t-butoxycarbonyl-3-pyrrolin-3-yl phenyl sulfone (4.46 g) in the same manner as described for REFERENCE EXAMPLE 1.

MS (CI<sup>+</sup>) *m/z*: 285 (MH<sup>+</sup>).

HRMS (CI<sup>+</sup>) for C<sub>17</sub>H<sub>21</sub>N<sub>2</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 285.1603; found, 285.1616.

#### Step 2.

1-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-Cyano-3-trifluoroacetyl-3-azabicyclo[3.1.0]hexan-6-yl)]-4-nitrobenzene.

To a solution of (1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)benzene (853 mg) in dichloromethane (7 mL) was added

trifluoroacetic acid (7 mL) at 0 °C, the mixture was stirred at room temperature for 75 minutes and concentrated in vacuo. To a solution of the residue in dichloromethane (7 mL) was added triethylamine (5.01 mL) and trifluoroacetic anhydride (1.06 mL) at 0 °C, the mixture was stirred at room temperature overnight and concentrated in vacuo. After dilution of the residue with ethyl acetate, the mixture was washed with 1 N hydrochloric acid, water, aqueous sodium hydrogencarbonate solution and brine, dried over anhydrous sodium sulfate, filtered, and then concentrated in vacuo. To a solution of the residue in chloroform (3 mL) was added ammonium nitrate (372 mg) and trifluoroacetic anhydride (2.19 mL), the mixture was stirred at room temperature for 2.7 hours. After addition of ice, the resulting precipitates were collected by filtration and washed with water and dichloromethane to give 1-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-trifluoroacetyl-3-azabicyclo[3.1.0]hexan-6-yl)]-4-nitrobenzene (546 mg). MS (EI<sup>+</sup>) *m/z*: 325 (M<sup>+</sup>).

HRMS (EI<sup>+</sup>) for C<sub>14</sub>H<sub>10</sub>F<sub>3</sub>N<sub>3</sub>O<sub>3</sub> (M<sup>+</sup>): calcd, 325.0674; found, 325.0648.

### 15 Step 3.

1-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-*t*-Butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]-4-nitrobenzene.

The mixture of 1-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-trifluoroacetyl-3-azabicyclo[3.1.0]hexan-6-yl)]-4-nitrobenzene (9.2 mg) and a solution of ammonia in methanol (6.7 M, 0.5 mL) was stirred at room temperature for 21 hours and concentrated in vacuo. To a solution of the residue in tetrahydrofuran (0.5 mL) was added triethylamine (19.7  $\mu$ L) and di-*t*-butyl dicarbonate (9.5 mg) at 0 °C, the mixture was stirred at room temperature for 30 minutes. Preparative thin layer chromatography (silica, hexane : ethyl acetate = 4:5) of the residue gave 1-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-*t*-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]-4-nitrobenzene (8.7 mg).

MS (EI<sup>+</sup>) *m/z*: 329 (M<sup>+</sup>).

HRMS (EI<sup>+</sup>) for C<sub>17</sub>H<sub>19</sub>N<sub>3</sub>O<sub>4</sub> (M<sup>+</sup>): calcd, 329.1376; found, 329.1401.

### 30 REFERENCE EXAMPLE 4

1-Benzyloxycarbonylamino-4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-*t*-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]benzene.

The title compound 1-benzyloxycarbonylamino-4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-*t*-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]benzene (127 mg) was prepared from 1-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-*t*-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-

6-yl)]-4-nitrobenzene (98.8 mg) in the same manner as described for REFERENCE EXAMPLE 1.

MS (EI<sup>+</sup>) *m/z*: 433 (M<sup>+</sup>).

5 HRMS (EI<sup>+</sup>) for C<sub>25</sub>H<sub>27</sub>N<sub>3</sub>O<sub>4</sub> (M<sup>+</sup>): calcd, 433.2002; found, 433.1989.

#### Antibacterial Activity

The pharmaceutically-acceptable compounds of the present invention are useful antibacterial agents having a good spectrum of activity in vitro against standard bacterial strains, which are used to screen for activity against pathogenic bacteria. Notably, the pharmaceutically-acceptable compounds of the present invention show activity against vancomycin-resistant enterococci, streptococci including penicillin-resistant *S. pneumoniae*, methicillin-resistant *S. aureus*, *M. catarrhalis*, and *C. pneumoniae*. The antibacterial spectrum and potency of a particular compound may be determined in a standard test system.

15 The following in vitro results were obtained based on an agar dilution method except for *C. pneumoniae*. The activity is presented as the minimum inhibitory concentration (MIC).

*S. aureus* and *M. catarrhalis* were tested on Mueller-Hinton agar, using an approximate inoculum of 1 x 10<sup>4</sup> cfu/spot an incubation temperature of 35°C for 24 hours. The MIC was defined as the lowest concentration at which no visible bacterial growth was observed.

Streptococci and enterococci were tested on Mueller-Hinton agar supplemented with 5 % defibrinated horse blood, using an approximate inoculum of 1 x 10<sup>4</sup> cfu/spot an incubation temperature of 35°C in an atmosphere of 5 % CO<sub>2</sub> for 24 hours. The MIC was defined as the lowest concentration at which no visible bacterial growth was observed.

*C. pneumoniae* was tested using minimum essential medium supplemented with 10 % heat-inactivated fetal bovine serum, 2 mM L-glutamine, 1 mg/ml cycloheximide and non essential amino acid. HeLa 229 cells were inoculated with 10<sup>4</sup> inclusion-forming units of *C. pneumoniae* strain per mL. Infected cells were incubated with test compounds in complete medium at 35°C in an atmosphere of 5 % CO<sub>2</sub> for 72 hours. Cells monolayers were fixed in methanol, stained for chlamydial inclusions with a fluorescein-conjugated anti-Chlamydia monoclonal antibody, and were observed with fluorescence microscope. The MIC was defined as the lowest concentration at which no inclusion was observed.

35


Strains	MIC ( $\mu\text{g/ml}$ )				
	example 1	example 8	example 9	example 11	Linezolid
<i>Staphylococcus aureus</i>					
Smith	0.25	1	0.5	0.25	1
CR	2	2	1	1	16
MR	0.25	1	0.5	0.5	1
<i>Streptococcus pneumoniae</i>					
IID553	0.5	0.5	0.5	0.25	2
PRQR	0.25	0.5	0.5	0.25	1
<i>Streptococcus pyogenes</i>					
IID692	0.5	0.5	0.5	0.125	1
<i>Enterococcus faecium</i>					
VRQR	0.25	0.5	0.5	0.25	2
<i>Moraxella catarrhalis</i>					
ATCC25238	0.5	2	2	1	4

- CR = chloramphenicol resistant!  
MR = methicillin resistant  
PRQR = penicillin resistant, quinolone resistant  
5 VRQR = vancomycin resistant, quinolone resistant  
NT = not tested

The invention described herein is exemplified by the following non-limiting examples. The compound data is designated in accordance to *General Guidelines for*  
10 *Manuscript Preparation*, J. Org. Chem. Vol. 66, pg. 19A, Issue 1, 2001.

## WHAT IS CLAIMED IS:

1. The present invention relates to compounds of formula I:



I

its enantiomer, diastereomer, or pharmaceutically acceptable salt, hydrate or prodrug thereof  
wherein:

R<sub>1</sub> represents

vi) hydrogen,

vii) NR<sub>5</sub>R<sub>6</sub>,

15 viii) CR<sub>7</sub>R<sub>8</sub>R<sub>9</sub>, C(R)<sub>2</sub>OR<sub>14</sub>, CH<sub>2</sub>NHR<sub>14</sub>,

ix) C(=O)R<sub>13</sub>, C(=NOH)H, C(=NOR<sub>13</sub>)H, C(=NOR<sub>13</sub>)R<sub>13</sub>, C(=NOH)R<sub>13</sub>, C(=O)N(R<sub>13</sub>)<sub>2</sub>,  
C(=NOH)N(R<sub>13</sub>)<sub>2</sub>, NH<sub>2</sub>C(=X<sub>1</sub>)N(R<sub>13</sub>)<sub>2</sub>, (C=NH)R<sub>7</sub>, N(R<sub>13</sub>)C(=X<sub>1</sub>)N(R<sub>13</sub>)<sub>2</sub>, COOR<sub>13</sub>,  
SO<sub>2</sub>R<sub>14</sub>, N(R<sub>13</sub>)SO<sub>2</sub>R<sub>14</sub>, N(R<sub>13</sub>)COR<sub>14</sub>,

x) (C<sub>1-6</sub>alkyl)CN, CN, CH=C(R)<sub>2</sub>, (CH<sub>2</sub>)<sub>p</sub>OH, C(=O)CHR<sub>13</sub>, C(=NR<sub>13</sub>)R<sub>13</sub>,

20 NR<sub>10</sub>C(=X<sub>1</sub>)R<sub>13</sub>; or

vi) C<sub>5-10</sub> heterocycle optionally substituted with 1-3 groups of R<sub>7</sub>, which may be attached  
through either a carbon or a heteroatom;

25 A represents NR, O, or S(O)<sub>p</sub>;





represents aryl or heteroaryl, heterocycle, heterocyclyl or heterocyclic, provided that in the case of a heteroaryl, heterocycle, heterocyclyl or heterocyclic, the cyclopropyl is not attached to a nitrogen atom on the ring;

5  $R_x$  represents hydrogen or C<sub>1-6</sub> alkyl;

$R_3$  represent

- i)  $NR_{13}(C=X_2)R_{12}$ ,
- ii)  $NR_{13}(C=X_1)R_{12}$ ,
- 10 iii)  $NR_{13}SO_2R_{14}$ ,
- iv)  $N(R_{13})\text{heteroaryl}$ ,
- v)  $NR_{13}(CHR_{13})_{0-4}\text{aryl}$ ,
- vi)  $NR_{13}(CHR_{13})_{0-4}\text{heteroaryl}$ ,
- vii)  $S(CHR_{13})_{0-4}\text{aryl}$ ,
- 15 viii)  $S(CHR_{13})_{0-4}\text{heteroaryl}$ ,
- ix)  $O(CHR_{13})_{0-4}\text{aryl}$ ,
- x)  $O(CHR_{13})_{0-4}\text{heteroaryl}$ ,
- xi)  $NOH(C=X_1)R_{12}$ ,
- xii)  $-OC=N(OCO\text{aryl})$  C<sub>1-6</sub> alkyl
- 20 xiii)  $-OC=N(\overset{\cdot}{O}H)C_{1-6}$  alkyl
- xiv) C<sub>5-10</sub> heteroaryl which may be attached through either a carbon or a heteroatom; said aryl and heteroaryl optionally substituted with 1-3 groups of  $R_7$ ,

$R_4$ , and  $R_{4a}$ , independently represent

- 25 v) hydrogen,
  - vi) halogen,
  - vii) C<sub>1-6</sub> alkoxy, or
  - viii) C<sub>1-6</sub> alkyl
- 30  $r$  and  $s$  independently are 1-3, with the provision that when  $(R_{4a})_s$  and  $(R_4)_r$  are attached to an Ar or HAr ring the sum of  $r$  and  $s$  is less than or equal to 4;

R<sub>5</sub> and R<sub>6</sub> independently represent

- xiii) hydrogen,
- xiv) C<sub>1-6</sub> alkyl optionally substituted with 1-3 groups of halogen, CN, OH, C<sub>1-6</sub> alkoxy, amino, imino, hydroxyamino, alkoxyamino, C<sub>1-6</sub> acyloxy, C<sub>1-6</sub> alkylsulfenyl, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylsulfonyl, aminosulfonyl, C<sub>1-6</sub> alkylaminosulfonyl, C<sub>1-6</sub> dialkylaminosulfonyl, 4-morpholinylsulfonyl, phenyl, pyridine, 5-isoxazolyl, ethylenyloxy, or ethynyl, said phenyl and pyridine optionally substituted with 1-3 halogen, CN, OH, CF<sub>3</sub>, C<sub>1-6</sub> alkyl or C<sub>1-6</sub> alkoxy;
- 5      xv) C<sub>1-6</sub> acyl optionally substituted with 1-3 groups of halogen, OH, SH, C<sub>1-6</sub> alkoxy, naphthalenoxy, phenoxy, amino, C<sub>1-6</sub> acylamino, hydroxylamino, alkoxyamino, C<sub>1-6</sub> acyloxy, aralkyloxy, phenyl, pyridine, C<sub>1-6</sub> alkylcarbonyl, C<sub>1-6</sub> alkylamino, C<sub>1-6</sub> dialkylamino, C<sub>1-6</sub> hydroxyacyloxy, C<sub>1-6</sub> alkylsulfenyl, phthalimido, maleimido, succinimido, said phenoxy, phenyl and pyridine optionally substituted with 1-3 groups of halo, OH, CN, C<sub>1-6</sub> alkoxy, amino, C<sub>1-6</sub> acylamino, CF<sub>3</sub> or C<sub>1-6</sub> alkyl;
- 10      xvi) C<sub>1-6</sub> alkylsulfonyl optionally substituted with 1-3 groups of halogen, OH, C<sub>1-6</sub> alkoxy, amino, hydroxylamino, alkoxyamino, C<sub>1-6</sub> acyloxy, or phenyl; said phenyl optionally substituted with 1-3 groups of halo, OH, C<sub>1-6</sub> alkoxy, amino, C<sub>1-6</sub> acylamino, CF<sub>3</sub> or C<sub>1-6</sub> alkyl;
- 15      xvii) arylsulfonyl optionally substituted with 1-3 of halogen, C<sub>1-6</sub> alkoxy, OH or C<sub>1-6</sub> alkyl;
- 20      xviii) C<sub>1-6</sub> alkoxy carbonyl optionally substituted with 1-3 of halogen, OH, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> acyloxy, or phenyl, said phenyl optionally substituted with 1-3 groups of halo, OH, C<sub>1-6</sub> alkoxy, amino, C<sub>1-6</sub> acylamino, CF<sub>3</sub> or C<sub>1-6</sub> alkyl;
- 25      xix) aminocarbonyl, C<sub>1-6</sub> alkylaminocarbonyl or C<sub>1-6</sub> dialkylaminocarbonyl, said alkyl groups optionally substituted with 1-3 groups of halogen, OH, C<sub>1-6</sub> alkoxy or phenyl
- 20      xx) five to six membered heterocycles optionally substituted with 1-3 groups of halogen, OH, CN, amino, C<sub>1-6</sub> acylamino, C<sub>1-6</sub> alkylsulfonylamino, C<sub>1-6</sub> alkoxy carbonylamino, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> acyloxy or C<sub>1-6</sub> alkyl, said alkyl optionally substituted with 1-3 groups of halogen, or C<sub>1-6</sub> alkoxy;
- 30      xxi) C<sub>3-6</sub> cycloalkylcarbonyl optionally substituted with 1-3 groups of halogen, OH, C<sub>1-6</sub> alkoxy or CN;
- xxii) benzoyl optionally substituted with 1-3 groups of halogen, OH, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkyl, CF<sub>3</sub>, C<sub>1-6</sub> alkanoyl, amino or C<sub>1-6</sub> acylamino;
- xxiii) pyrrolylcarbonyl optionally substituted with 1-3 of C<sub>1-6</sub> alkyl;

xxiv) C<sub>1-2</sub> acyloxyacetyl where the acyl is optionally substituted with amino, C<sub>1-6</sub> alkylamino, C<sub>1-6</sub> dialkylamino, 4-morpholino, 4-aminophenyl, 4-(dialkylamino)phenyl, 4-(glycylamino)phenyl; or

R<sub>5</sub> and R<sub>6</sub> taken together with any intervening atoms can form a 3 to 7 membered  
 5 heterocyclic ring containing carbon atoms and 1-2 heteroatoms independently chosen from O, S, SO, SO<sub>2</sub>, N, or NR<sub>8</sub>;

R<sub>7</sub> represent

iii) hydrogen, halogen, CN, CO<sub>2</sub>R, CON(R)<sub>2</sub>, CHO, CH<sub>2</sub>NHAc, C(=NOR), OH, C<sub>1-6</sub>  
 10 alkoxy, C<sub>1-6</sub> alkyl, alkenyl, hydroxy C<sub>1-6</sub> alkyl, (CH<sub>2</sub>)<sub>1-3</sub>NHC(O)C<sub>1-6</sub> alkyl, (CH<sub>2</sub>)<sub>1-3</sub>N(C<sub>1-6</sub> alkyl)<sub>2</sub>

iv) (CH<sub>2</sub>)<sub>n</sub>amino, (CH<sub>2</sub>)<sub>n</sub>C<sub>1-6</sub> alkylamino, C<sub>1-6</sub> dialkylamino, hydroxylamino or C<sub>1-2</sub>  
 alkoxyamino all of which can be optionally substituted on the nitrogen with C<sub>1-6</sub>  
 acyl, C<sub>1-6</sub> alkylsulfonyl or C<sub>1-6</sub> alkoxycarbonyl, said acyl and alkylsulfonyl  
 15 optionally substituted with 1-2 of halogen or OH;

R<sub>8</sub> and R<sub>9</sub> independently represents

iv) H, CN,  
 v) C<sub>1-6</sub> alkyl optionally substituted with 1-3 halogen, CN, OH, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub>  
 20 acyloxy, or amino,  
 vi) phenyl optionally substituted with 1-3 groups of halogen, OH, C<sub>1-6</sub> alkoxy; or

R<sub>7</sub> and R<sub>8</sub> taken together can form a 3-7 membered carbon ring optionally interrupted with  
 1-2 heteroatoms chosen from O, S, SO, SO<sub>2</sub>, NH, and NR<sub>8</sub>;

25 X<sub>1</sub> represents O, S or NR<sub>13</sub>, NCN, NCO<sub>2</sub>R<sub>16</sub>, or NSO<sub>2</sub>R<sub>14</sub>

X<sub>2</sub> represents O, S, NH or NSO<sub>2</sub>R<sub>14</sub>;

30 R<sub>10</sub> represents hydrogen, C<sub>1-6</sub> alkyl or CO<sub>2</sub>R<sub>15</sub>;

R<sub>12</sub> represents hydrogen, C<sub>1-6</sub> alkyl, NH<sub>2</sub>, OR, CHF<sub>2</sub>, CHCl<sub>2</sub>, CR<sub>2</sub>Cl, (CH<sub>2</sub>)<sub>n</sub>SR, (CH<sub>2</sub>)<sub>n</sub>CN, (CH<sub>2</sub>)<sub>n</sub>SO<sub>2</sub>R, (CH<sub>2</sub>)<sub>n</sub>S(O)R, C<sub>1-6</sub> alkylamino, C<sub>5-10</sub> heteroaryl or C<sub>1-6</sub> dialkylamino, where

said alkyl may be substituted with 1-3 groups of halo, CN, OH or C<sub>1-6</sub> alkoxy, said heteroaryl optionally substituted with 1-3 groups of R<sub>7</sub>;

Each R<sub>13</sub> represents independently hydrogen, C<sub>1-6</sub> alkyl, C<sub>6-10</sub> aryl, NR<sub>5</sub>R<sub>6</sub>, SR<sub>8</sub>, S(O)R<sub>8</sub>, S(O)<sub>2</sub>R<sub>8</sub>, CN, OH, C<sub>1-6</sub> alkylS(O)R, C<sub>1-6</sub> alkoxycarbonyl, hydroxycarbonyl, -OCOaryl, C<sub>1-6</sub> acyl, C<sub>3-7</sub> membered carbon ring optionally interrupted with 1-4 heteroatoms chosen from O, S, SO, SO<sub>2</sub>, NH and NR<sub>8</sub> where said C<sub>1-6</sub> alkyl, aryl or C<sub>1-6</sub> acyl groups may be independently substituted with 0-3 halogens, hydroxy, N(R)<sub>2</sub>, CO<sub>2</sub>R, C<sub>6-10</sub> aryl, C<sub>5-10</sub> heteroaryl, or C<sub>1-6</sub> alkoxy groups;

When two R<sub>13</sub> groups are attached to the same atom or two adjacent atoms they may be taken together to form a 3-7 membered carbon ring optionally interrupted with 1-2 heteroatoms chosen from O, S, SO, SO<sub>2</sub>, NH, and NR<sub>8</sub>;

R represents hydrogen or C<sub>1-6</sub> alkyl;

R<sub>14</sub> represents amino, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, five to six membered heterocycles or phenyl, said phenyl and heterocycles optionally substituted with 1-3 group of halo, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> acylamino, or C<sub>1-6</sub> alkyl, hydroxy and/or amino, said amino and hydroxy optionally protected with an amino or hydroxy protecting group;

R<sub>15</sub> is C<sub>1-6</sub> alkyl or benzyl said benzyl optionally substituted with 1-3 groups of halo, OH, C<sub>1-6</sub> alkoxy, amino, C<sub>1-6</sub> acylamino, or C<sub>1-6</sub> alkyl;


R<sub>16</sub> is hydrogen, C<sub>5-10</sub> heteroaryl, C<sub>6-10</sub> aryl, said heteroaryl and aryl optionally substituted with 1-3 groups of R<sub>7</sub>;

p represents 0-2 and

m, n, and q represents 0-1.

30

2. A compound according to claim 1 wherein R<sub>1</sub> represents H, NR<sub>5</sub>R<sub>6</sub>, CN, OH, C(R)<sub>2</sub>OR<sub>14</sub>, NHC(=X<sub>1</sub>)N(R<sub>13</sub>)<sub>2</sub>, C(=NOH)N(R<sub>13</sub>)<sub>2</sub>, NR<sub>10</sub>C(=X<sub>1</sub>)R<sub>13</sub> or CR<sub>7</sub>R<sub>8</sub>R<sub>9</sub>.




3. A compound according to claim 1 wherein  
is phenyl, pyridine, pyrimidine, or piperidine.

4. A compound according to claim 1 wherein  $R_1$  is  $NR_5R_6$ .

5. A compound according to claim 1 wherein  $R_1$  is CN.

6. A compound according to claim 1 wherein one of is  $NR_{10}C(=X_1)R_{13}$ .

7. A compound according to claim 1 wherein  $R_3$  is  $NR(C=X_1)R_{12}$ ,  
C5-10 heteroaryl,  $NH(CH_2)_{0-4}$ aryl,  $NH(CH_2)_{0-4}$ heteroaryl, said aryl and heteroaryl optionally  
substituted with 1-3 groups of  $R_a$ .

8. A compound according to claim 7 wherein  $R_3$  is a C5-10 heteroaryl  
represented by  which represents an optionally substituted aromatic heterocyclic group  
containing 1 to 4 nitrogen atoms and at least one double bond, and which is connected  
through a bond on any nitrogen.

9. A compound according to claim 1 wherein  $R_5$  and  $R_6$  independently  
are:


- v) H,
- vi) C<sub>1-6</sub> alkyl optionally substituted with 1-3 groups of halogen, CN, OH, C<sub>1-6</sub> alkoxy,  
amino, hydroxyamino, alkoxyamino, C<sub>1-6</sub> acyloxy, C<sub>1-6</sub> alkylsulfenyl, C<sub>1-6</sub>  
alkylsulfinyl, C<sub>1-6</sub> alkylsulfonyl, aminosulfonyl, C<sub>1-6</sub> alkylaminosulfonyl, C<sub>1-6</sub>  
dialkylaminosulfonyl, 4-morpholinylsulfonyl, phenyl, pyridine, 5-isoxazolyl,  
ethylenoxy, or ethynyl, said phenyl and pyridine optionally substituted with 1-3  
halogen, CN, OH, CF<sub>3</sub>, C<sub>1-6</sub> alkyl or C<sub>1-6</sub> alkoxy;
- vii) C<sub>1-6</sub> acyl optionally substituted with 1-3 groups of halogen, OH, SH, C<sub>1-6</sub> alkoxy,  
naphthalenoxy, phenoxy, amino, C<sub>1-6</sub> acylamino, hydroxylamino, alkoxylamino, C<sub>1-6</sub>  
acyloxy, phenyl, pyridine, C<sub>1-6</sub> alkylcarbonyl, C<sub>1-6</sub> alkylamino, C<sub>1-6</sub>  
dialkylamino, C<sub>1-6</sub> hydroxyacyloxy, C<sub>1-6</sub> alkylsulfenyl, phthalimido, maleimido,

- viii) succinimido, said phenoxy, phenyl and pyridine optionally substituted with 1-3 groups of halo, OH, CN, C<sub>1-6</sub> alkoxy, amino, C<sub>1-6</sub> acylamino, CF<sub>3</sub> or C<sub>1-6</sub> alkyl; or benzoyl optionally substituted with 1-3 groups of halogen, OH, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkyl, CF<sub>3</sub>, C<sub>1-6</sub> alkanoyl, amino or C<sub>1-6</sub> acylamino and all other variables are as described herein.

10. A compound according to claim 1 wherein X<sub>1</sub> represents O.

11. A compound according to claim 1 wherein the structural formula is

10 II:



15

Formula II

wherein R<sub>1</sub>, R<sub>4</sub>, R<sub>4a</sub>, Y and R<sub>3</sub> are as described above.

12. A compound which is:
- 20 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyanobicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 25 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride,  
 1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole,

- 1-[5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]-1,2,3-triazole hydrochloride,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-acetoxyacetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 5 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-hydroxyacetyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-methanesulfonyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-methyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 10 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3,6-dicyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-cyanomethyl-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 15 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[(isoxazol-3-yl)oxy]methyloxazolidin-2-one,  
 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[(isoxazol-3-yl)oxy]methyloxazolidin-2-one hydrochloride,  
 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(3-t-butoxycarbonyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[N-(t-butoxycarbonyl)-N-(isoxazol-3-yl)]aminomethyloxazolidin-2-one,  
 20 5(R)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-(6-cyano-3-azabicyclo[3.1.0]hexan-6-yl)]phenyl]-5-[N-(isoxazol-3-yl)]aminomethyloxazolidin-2-one hydrochloride,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(5-cyanopyridin-2-yl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 25 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(pyridin-2-yl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-acetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(pyrimidin-2-yl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 30 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(4-pyridylmethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(N-cyano-1-iminoethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,

- N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-methoxycarbonyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(N-cyano-S-methylthioiminomethyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 5 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(N-cyanocarboxamidyl)-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-(N,N'-t-butoxycarbonylcarboxamidyl)-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-carboxamidyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-  
 10 2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-(N-t-Butoxycarbonylamino)acetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[3-aminoacetyl-6-cyano-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-  
 2-oxooxazolidin-5-ylmethyl]acetamide hydrochloride,  
 15 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-methanesulfonylacetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(dibenzylphosphoryloxy)acetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 N-[5(S)-3-[4-[(1 $\alpha$ ,5 $\alpha$ ,6 $\beta$ )-[6-cyano-3-(phosphoryloxy)acetyl-3-azabicyclo[3.1.0]hexan-6-yl]]phenyl]-2-oxooxazolidin-5-ylmethyl]acetamide,  
 20

or its enantiomer, diastereomer, or pharmaceutically acceptable salt, hydrate or prodrug thereof.

- 25                    13.     A pharmaceutical composition comprised of a compound in accordance with claim 1 in combination with a pharmaceutically acceptable carrier.
14.     A pharmaceutical composition produced by combining a compound in accordance with claim 1 with a pharmaceutically acceptable carrier.
- 30                    15.     A method of treating or preventing a bacterial infection in a mammalian patient in need thereof, comprising administering to said patient an effective amount of a compound of claim 1.



16. A method of treating or preventing bacterial infection by administering an effective amount of a compound of formula I of claim 1 and an effective amount of one or more of a vitamin selected from the group consisting of vitamin B2, vitamin B6, vitamin B12 and folic acid to a patient in need thereof.

5

17. A method of treating or preventing an oxazolidinone-associated side effect by administering an effective amount of the oxazolidinone of structural formula I of claim 1 and an effective amount of one or more of vitamin B2, vitamin B6, vitamin B12 and folic acid to a patient in need thereof.

10

18. A method according to claim 17 for treating or preventing oxazolidinone-associated normocytic anemia or peripheral sensory neuropathy by administering an effective amount of vitamin B2 to a patient in need thereof.

15

19. A method according to claim 17 for treating or preventing oxazolidinone-associated sideroblastic anemia, peripheral sensory neuropathy, optic neuropathy, seizures, thrombocytopenia, cheilosis, and seborrheic dermatitis by administering an effective amount of vitamin B6 to a patient in need thereof.

20

20. A method according to claim 17 for treating or preventing oxazolidinone-associated hypo-regenerative anemia, megaloblastic anemia by administering an effective amount of vitamin B12 and folic acid to a patient in need thereof.

25

21. A pharmaceutical composition comprised of a compound in accordance with claim 1 in combination with a vitamin selected from the group consisting of vitamin B2, vitamin B6, vitamin B12 and folic acid and a pharmaceutically acceptable carrier.

ABSTRACT OF THE DISCLOSURETITLE OF THE INVENTION


## Oxazolidinone Antibiotics and Derivatives Thereof

5

This invention relates to new oxazolidinones having a cyclopropyl moiety, which are effective against aerobic and anerobic pathogens such as multi-resistant staphylococci, streptococci and enterococci, Bacteroides spp., Clostridia spp. species, as well as acid-fast organisms such as *Mycobacterium tuberculosis* and other mycobacterial species.

10

The compounds are represented by structural formula I:



I

15 its enantiomer, diastereomer, or pharmaceutically acceptable salt or ester thereof.

20

WDC99 894402-1.050193.0178

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/020736

International filing date: 29 June 2004 (29.06.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/553,963  
Filing date: 18 March 2004 (18.03.2004)

Date of receipt at the International Bureau: 22 September 2004 (22.09.2004)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse